

Representations of $\mathrm{PGL}(2)$ of a local field and harmonic forms on simplicial complexes

Paul Broussous
 UMR6086 CNRS
 SP2MI - Téléport 2
 Bd M. et P. Curie BP 30179
 86962 Futuroscope Chasseneuil Cedex
 E-mail : broussou@math.univ-poitiers.fr

January 2007

Dedicated to Colin Bushnell for his 60th birthday

Abstract

Nous donnons des modèles combinatoires des représentations lisses, complexes, génériques, non-sphériques du groupe $G = \mathrm{PGL}(2, F)$, où F est un corps localement compact non-archimédien. Plus précisément nous reprenons l'étude des graphes $(\tilde{X}_k)_{k \geq 0}$ inaugurée dans un précédent travail. Nous montrons que de telles représentations se réalisent comme quotients de la cohomologie d'un graphe $\tilde{X}_k$ pour un k bien choisi, ou, de façon équivalente, dans un espace de formes harmoniques discrètes sur un tel graphe. Pour les représentations supercuspidales, ces modèles sont de plus uniques.

Introduction

Let F be a non-archimedean local field and G be the locally compact group $\mathrm{PGL}(N, F)$, where $N \geq 2$ is an integer. In [1] the author constructed a projective tower of simplicial complexes fibered over the Bruhat-Tits building X of G . He addressed the question of determining the structure of the cohomology spaces of these complexes as G -modules. In this article we give a conceptual treatment of the case

$N = 2$. In that case X is a homogeneous tree and (a slightly modified version of) the projective tower is formed of graphs $\tilde{X}_n$, $n \geq 0$, acted upon by G .

Let π be an irreducible smooth complex representation of G that we assume generic and non-spherical. We show there is a natural G -equivariant map

$$\tilde{\Psi}_\pi : \tilde{\pi} \longrightarrow \mathcal{H}_\infty(\tilde{X}_{n(\pi)}, \mathbb{C}) ,$$

where $n(\pi)$ is an integer related to the conductor of π , $\tilde{\pi}$ denotes the contragredient representation of π , and $\mathcal{H}_\infty(\tilde{X}_{n(\pi)}, \mathbb{C})$ denotes the space of smooth discrete harmonic forms on $\tilde{X}_{n(\pi)}$. Our construction is based on the existence of *new vectors* for irreducible generic representations whose proof is due to Casselman in the case $N = 2$ [4] (see [6] for the general case).

The G -space $\mathcal{H}_\infty(\tilde{X}_{n(\pi)}, \mathbb{C})$ is naturally isomorphic to the contragredient representation of the cohomology space $H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$ (cohomology space with compact support and complex coefficients). We show that $\tilde{\Psi}_\pi$ corresponds to a natural G -equivariant map:

$$\Psi_\pi : H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C}) \longrightarrow \pi .$$

In other words π is naturally a quotient of $H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$. When π is supercuspidal the surjective map Ψ splits and π embeds in $H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$. We show that this model is unique:

$$\dim_{\mathbb{C}} \text{Hom}_G(\pi, H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})) = 1 .$$

The proof of that fact roughly goes as follows. Using a *geodesic Radon transform* on the space of 1-cochains with finite support on $\tilde{X}_k$, $k \geq 0$, we construct an intertwining operator:

$$j_k : H_c^1(\tilde{X}_k, \mathbb{C}) \longrightarrow c\text{-ind}_T^G 1_T ,$$

(the compactly induced representation of the trivial character of the diagonal torus T of G). The point is that this map is injective. Moreover we have:

$$\bigcup_{k \geq 0} \text{Im}(j_k) = c\text{-ind}_T^G 1_T .$$

We show that $c\text{-ind}_T^G 1_T$ naturally embeds in a space of Whittaker functions on G and we may then rely on the uniqueness of Whittaker model for $\text{PGL}(2, F)$.

That such a combinatorial realization of the generic non-spherical irreducible representations of $\mathrm{PGL}(2, F)$ is feasible was actually conjectured by Pierre Cartier more than thirty years ago [3] (but he did not introduce any simplicial structure). Later on Cartier's student Ahumada Bustamante [2] studied the action of the full automorphism group $\Gamma = \mathrm{Aut}(X)$ of the tree X on pairs of vertices at distance $k + 1$ (i.e. on edges of $\tilde{X}_k$). Using an equivalent language, he proved that, under the action of Γ , the space $\mathcal{H}_2(\tilde{X}_k, \mathbb{C})$ of L^2 harmonic forms splits into two irreducible components $\mathcal{H}_2^\pm(\tilde{X}_k, \mathbb{C})$, formed of even and odd harmonic forms respectively.

I would like to thank V. Sécherre and A. Bouaziz for pointing out some embarrassing mistakes in previous versions of this work. I thank G. Henniart, A. Gaborieau and A. Raghuram for stimulating discussions.

The notes are organized as follows. In §1 we shall discuss the link between cohomology and harmonic forms on graphs. §2,3,5 are about the construction of the maps Ψ_π , $\tilde{\Psi}_\pi$ and some of their properties. The Radon transform is defined and studied in §5 and is used in §6 to prove our uniqueness result.

In the sequel we shall use the following notation:

$\mathfrak{o} = \mathfrak{o}_F$ is the ring of integers of F ,
 $\mathfrak{p} = \mathfrak{p}_F$ is the maximal ideal of $\mathfrak{o}$,
 $v = v_F$ is the normalized additive valuation of F ,
 $k = k_F$ is the residue class field $\mathfrak{o}/\mathfrak{p}$,
 $q = |k_F|$ is the cardinal of k ,
 $|| = ||_F$ is the multiplicative valuation on $F^\times$ normalized in such a way that $|\varpi|_F = 1/q$ for any generator ϖ of the ideal $\mathfrak{p}$.

1. Proper G -graphs and harmonic forms

1.1. In this section we let G be any locally profinite group and Y be a locally finite directed graph (each vertex belongs to a finite number of edges). We write Y^0 (resp. Y^1) for the set of vertices (resp. edges) of Y . We have the map $Y^1 \longrightarrow Y^0$, $a \mapsto a^+$ (resp. $a \mapsto a^-$), where for any edge a we denote by a^+ and a^- its head and tail respectively. We assume that G acts on Y by preserving the structure of directed graph. For all $s \in Y^0$, $a \in Y^1$, we have incidence numbers $[a : s] \in \{-1, 1, 0\}$ satisfying $[g.a : g.s] = [a : s]$, for all $g \in G$; there are defined by

$[a : a^+] = +1$, $[a : a^-] = -1$, and $[a : s] = 0$ if $s \notin \{a^+, a^-\}$. Finally we assume that the action of G on Y is proper: for all $a \in Y^0$, the stabilizer $G_s := \{g \in G ; g.s = s\}$ is open and compact.

1.2. We let $H_c^1(Y, \mathbb{C})$ denote the cohomology space of the CW-complex Y with compact support and complex coefficients. Recall that it may be calculated as follows. Let $C_0(Y, \mathbb{C})$ (resp. $C_1(Y, \mathbb{C})$) be the $\mathbb{C}$ -vector space with basis Y^0 (resp. Y^1). Let $C_c^i(Y, \mathbb{C})$, $i = 1, 2$, be the $\mathbb{C}$ -vector space of 1-cochains with finite support : $C_c^i(Y, \mathbb{C})$ is the subspace of the algebraic dual of $C_i(Y, \mathbb{C})$ formed of those linear forms whose restrictions to the basis Y^i have finite support. The coboundary map

$$d : C_c^0(Y, \mathbb{C}) \longrightarrow C_c^1(Y, \mathbb{C})$$

is given by $df(a) = f(a^+) - f(a^-)$. Then

$$(1.2.1) \quad H_c^1(Y, \mathbb{C}) \simeq C_c^1(Y, \mathbb{C}) / dC_c^0(Y, \mathbb{C}) .$$

The group G acts on $C_i(Y, \mathbb{C})$ and $C_c^i(Y, \mathbb{C})$. Since the action of G on Y is proper, these spaces are smooth G -modules. The coboundary map is G -equivariant and the isomorphism (1.2.1) is G -equivariant. So $H_c^1(Y, \mathbb{C})$ is smooth as a G -module; it is not admissible in general.

1.3. For $i = 0, 1$, we have a natural pairing:

$$\langle -, - \rangle : C^i(Y, \mathbb{C}) \times C_c^i(Y, \mathbb{C}) \longrightarrow \mathbb{C} ,$$

where $C^i(Y, \mathbb{C})$ is the space of i -cochains with arbitrary support. The pairings are given by:

$$\langle f, g \rangle = \sum_{x \in Y^i} f(x)g(x) , \quad i = 0, 1 .$$

Via these pairings we may identify the algebraic dual $C_c^i(Y, \mathbb{C})^*$ of $C_c^i(Y, \mathbb{C})$ with $C^i(Y, \mathbb{C})$, $i = 0, 1$. The contragredient representation $C_c^i(Y, \mathbb{C})^\vee$ identifies with the space of smooth linear forms in $C^i(Y, \mathbb{C})$. A straightforward computation gives:

$$(1.3.1) \quad \langle f, dg \rangle = \langle d^*f, g \rangle , \quad f \in C^1(Y, \mathbb{C}) , \quad g \in C_c^0(Y, \mathbb{C}) ,$$

where $d^* : C^1(Y, \mathbb{C}) \longrightarrow C^0(Y, \mathbb{C})$ is defined by

$$d^*f(s) = \sum_{a \in Y^1, s \in a} [a : s]f(a) , \quad s \in Y^0 .$$

Of course this latter sum has a finite number of terms. An element of the kernel of d^* is called a *harmonic form* on Y . We denote by $\mathcal{H}(Y, \mathbb{C}) = \ker(d^*)$ this space of harmonic forms. It is naturally acted upon by G . The smooth part of $\mathcal{H}(Y, \mathbb{C})$ under the action of G , i.e. the space of *smooth harmonic forms* is denoted by $\mathcal{H}_\infty(Y, \mathbb{C})$. The following lemma follows from equality (1.3.1).

(1.3.2) **Lemma.** *The algebraic dual of $H_c^1(Y, \mathbb{C})$ naturally identifies with $\mathcal{H}(Y, \mathbb{C})$. Under this isomorphism, the contragredient representation of $H_c^1(Y, \mathbb{C})$ corresponds to $\mathcal{H}_\infty(Y, \mathbb{C})$.*

2. The projective tower of graphs

2.1 In this section, we recall the construction of [1]. The notation is slightly modified. We denote by X the Bruhat-Tits building of G (cf. [7] chap. II, §1). This is a 1-dimensional simplicial complex (a $(q+1)$ -homogeneous tree). Let $k \geq 0$ be an integer. An *(oriented) k -path* in X is an injective sequence $(s_0, \dots, s_k)$ of vertices in X such that, for $i = 0, \dots, k-1$, $\{s_i, s_{i+1}\}$ is an edge of X . We define an oriented graph $\tilde{X}_k$ as follows. Its vertex set (resp. edge set) is the set of k -paths (resp. $(k+1)$ -paths) in X . The structure of oriented graph is given by:

$$a^+ = (t_1, \dots, t_{k+1}), \quad a^- = (t_0, \dots, t_k), \quad \text{if } a = (t_0, \dots, t_{k+1}).$$

The group G acts on $\tilde{X}_k$. If $k \geq 1$, $\tilde{X}_k$ is a simplicial complex and the G -action is simplicial. For $k = 0$ the action preserves the graph structure. For all k , it preserves the orientation of $\tilde{X}_k$. Recall ([1] Lemma 4.1) that the simplicial complexes $\tilde{X}_k$, $k \geq 1$ are connected. The directed graph $\tilde{X}_0$ is obtained from X by doubling the edges (with the same vertex set); it is obviously connected.

2.2. For any integer $n \geq 1$, we write $\Gamma_0(\mathfrak{p}^n)$ for the image in G of the following subgroup of $\mathrm{GL}(2, F)$:

$$\tilde{\Gamma}_o(\mathfrak{p}^n) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}(2, F); \ a, d \in \mathfrak{O}^\times, \ b \in \mathfrak{O}, \ c \in \mathfrak{p}^n \right\}$$

We let $\Gamma_o(\mathfrak{p}^0)$ be the image in $\mathrm{PGL}(2, F)$ of the standard maximal compact subgroup of $\mathrm{GL}(2, F)$:

$$\tilde{\Gamma}_o(\mathfrak{p}^0) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}(2, \mathfrak{O}); \ ad - bc \in \mathfrak{O}^\times \right\}$$

For all $n \geq 0$, $\Gamma_o(\mathfrak{p}^{n+1})$ is the stabilizer in G of some edges of $\tilde{X}_n$. We fix such an edge a_o .

3. The construction

3.1. We start by recalling Casselman's result. We fix an irreducible complex smooth representation $(\pi, \mathcal{V})$ of G . We assume that:

(3.1.1) π has no non-zero vector fixed by $\Gamma_0(\mathfrak{p}^0)$,

(3.1.2) π is generic, i.e. it is not of the form $\chi \circ \det$, where χ is a character of $F^\times / ((F^\times)^2)$ and $\det : G \longrightarrow F^\times / (F^\times)^2$ is the map induced by the determinant map: $\mathrm{GL}(2, F) \longrightarrow F^\times$.

We have the following result ([4] Theorem 1):

(3.1.3) **Theorem** (Casselman). *i) For k large enough, the space of fixed vectors $\mathcal{V}^{\Gamma_o(\mathfrak{p}^{k+1})}$ is non-zero.*

ii) Let $n(\pi) \geq 0$ be such that $\mathcal{V}^{\Gamma_o(\mathfrak{p}^{n(\pi)+1})} \neq \{0\}$ and $\mathcal{V}^{\Gamma_o(\mathfrak{p}^{n(\pi)})} = \{0\}$. Then for all $k \geq n(\pi)$, we have:

$$\dim_{\mathbb{C}} \mathcal{V}^{\Gamma_o(\mathfrak{p}^{k+1})} = k - n(\pi) + 1 .$$

3.2. For all $a \in X_{n(\pi)}^1$ (resp. $s \in X_{n(\pi)}^0$), we write Γ_a (resp. Γ_s) for the stabilizer of a (resp. s) in G . In particular we have $\Gamma_{a_o} = \Gamma_o(\mathfrak{p}^{n(\pi)+1})$.

(3.2.1) **Lemma.** *i) For all $a \in X_{n(\pi)}^1$, we have $\dim \mathcal{V}^{\Gamma_a} = 1$.*

ii) Let $a \in X_{n(\pi)}^1$ and $s \in X_{n(\pi)}^0$ with $s \in a$. Then for all $v \in \mathcal{V}^{\Gamma_a}$, we have

$$\sum_{k \in \Gamma_s / \Gamma_a} kv = 0 .$$

Point i) is obvious. In ii), the vector $\sum_{k \in \Gamma_s / \Gamma_a} kv$ is fixed by Γ_s . So it must be zero since Γ_s is conjugate to $\Gamma_o(\mathfrak{p}^{n(\pi)})$.

Let us fix a non-zero vector $v_o \in \mathcal{V}^{\Gamma_{a_o}}$; v_o is unique up to a scalar in $\mathbb{C}^\times$. If a is any edge of $X_{n(\pi)}$, we put

$$(3.2.2) \quad v_a = gv_o, \text{ where } a = ga_o .$$

This is indeed possible since G acts transitively on $X_{n(\pi)}^1$. Moreover, since v_o is fixed by Γ_{a_o} , v_a does not depend on the choice of $g \in G$ such that $a = ga_o$. Let $\tilde{\mathcal{V}}$ be the contragredient representation of $\mathcal{V}$. We define a map:

$$\tilde{\Psi}_\pi : \tilde{\mathcal{V}} \longrightarrow C^1(X_{n(\pi)}, \mathbb{C})$$

by $\tilde{\Psi}_\pi(\varphi)(a) = \varphi(v_a)$. From (3.2.2) we have that $\tilde{\Psi}_\pi$ is G -equivariant.

(3.2.3) **Lemma.** *i) The image of $\tilde{\Psi}_\pi$ lies in $\mathcal{H}_\infty(X_{n(\pi)}, \mathbb{C})$.*

ii) The map $\tilde{\Psi}_\pi$ is injective.

It suffices to prove that $\text{Im}(\tilde{\Psi}_\pi) \subset \mathcal{H}(X_{n(\pi)}, \mathbb{C})$. So we must prove that for all $\varphi \in \tilde{\mathcal{V}}$, $(\varphi(v_a))_{a \in X_{n(\pi)}^1}$ is a harmonic form on $X_{n(\pi)}$, that is:

$$\sum_{s \in a} [a : s] \varphi(v_a) = 0, \text{ for all } s \in X_{n(\pi)}^0 .$$

Let s be any vertex of $X_{n(\pi)}$. Write $\bar{s}$ for the convex hull in X of the set vertices of X occuring in the path s (this is a segment lying in some apartment). We know that the pointwise stabilizer of $\bar{s}$ in G (that is the stabilizer Γ_s of s in G) acts transitively on the set of apartments of X containing $\bar{s}$. It follows that Γ_s acts transitively on

$$A_s^+ = \{a \in X_{n(\pi)}^1; a^+ = s\} \text{ and } A_s^- = \{a \in X_{n(\pi)}^1; a^- = s\} .$$

Fix some $a_s^+ \in A_s^+$ and $a_s^- \in A_s^-$. Then

$$\begin{aligned} \sum_{s \in a} [a : s] \varphi(v_a) &= \varphi \left(\sum_{a \in A_s^+} v_a - \sum_{a \in A_s^-} v_a \right) \\ &= \varphi \left(\sum_{k \in \Gamma_s / \Gamma_{a_s^+}} k v_{a_s^+} - \sum_{k \in \Gamma_s / \Gamma_{a_s^-}} k v_{a_s^-} \right) = 0 , \end{aligned}$$

thanks to lemma (3.2.1).

The G -equivariant map $\tilde{\Psi}_\pi$ is necessarily injective since it is non-zero and since the representation π is irreducible.

Passing to contragredient representations, we get an intertwining operator:

$$\tilde{\tilde{\Psi}}_\pi : \tilde{H}_c^1(\tilde{X}_{n(\pi)}, \mathbb{C}) \longrightarrow \tilde{\mathcal{V}} .$$

Recall that for any smooth G -module $\mathcal{W}$, we have a canonical injection $\mathcal{W} \longrightarrow \tilde{\mathcal{W}}$. It is surjective if and only if $\mathcal{W}$ is admissible. In particular $\mathcal{V}$ and $\tilde{\mathcal{V}}$ are canonically isomorphic since π is irreducible, whence admissible.

(3.2.4) **Theorem.** *The map $\tilde{\Psi}_\pi$ restricts to a non-zero intertwining operator $\Psi_\pi : H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C}) \longrightarrow \mathcal{V} \simeq \tilde{\mathcal{V}}$, given by:*

$$\Psi_\pi(\bar{\omega}) = \sum_{a \in \tilde{X}_{n(\pi)}^1} \omega(a) v_a .$$

where for $\omega \in C_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$, $\bar{\omega}$ denotes the image of ω in $H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$. In particular, the representation $(\pi, \mathcal{V})$ is naturally a subquotient of the cohomology space $H_c^1(X_{n(\pi)}, \mathbb{C})$.

The theorem follows from a straightforward computation based on Lemma (3.2.1)(ii) and is left to the reader.

Remark. Assume that π is supercuspidal. Then it is projective in the category of smooth complex representations of G . So as a corollary of Theorem (3.2.4) we get an injective map $(\pi, \mathcal{V}_\pi) \longrightarrow H_c^1(X_{n(\pi)}, \mathbb{C})$.

4. Some properties of the map $\tilde{\Psi}_\pi$.

4.1. We keep the notation as in the last section. If Y is any directed graph, we write $\mathcal{H}_c(Y, \mathbb{C})$ for the subspace of $\mathcal{H}(Y, \mathbb{C})$ of harmonic forms with finite support and $\mathcal{H}_2(X, \mathbb{C})$ for the subspace of L^2 -harmonic forms, that is forms $f \in \mathcal{H}(Y, \mathbb{C})$ satisfying

$$\sum_{a \in Y^1} |f(a)|^2 < \infty .$$

Note that any element of $\mathcal{H}_c(X_{n(\pi)}, \mathbb{C})$ is smooth.

(4.1.1) **Proposition.** *i) If π is a supercuspidal representation then $\text{Im} \tilde{P}si_\pi$ is contained in $\mathcal{H}_c(X_{n(\pi)}, \mathbb{C})$.*

ii) If π is a square-integrable representation then $\text{Im} \tilde{P}si_\pi$ lies in $\mathcal{H}_2(X_{n(\pi)}, \mathbb{C})$.

Assume π supercuspidal. Let $\lambda \in \tilde{\mathcal{V}}$. Then $\tilde{P}si_\pi(\lambda)(a) = \lambda(gv_{a_o})$ for all $a = ga_o \in X_{n(\pi)}^1$. Since π is supercuspidal, the coefficient

$g \in G \mapsto \lambda(gv_{a_o})$ has compact support C . Choose a finite number of compact open subgroups K_i , $i \in I$, of G and elements $g_i \in G$, $i \in I$, such that C lies in the union of the $g_i K_i$, $i \in I$. Then the support of the harmonic form $\tilde{P}si_\pi(\lambda)$ lies in

$$\bigcup_{i \in I} g_i K_i a_o = \bigcup_{i \in I} g_i K_i / (K_i \cap \Gamma_{a_o}) a_o ,$$

a finite set.

Now assume that π is square-integrable. With the notation as above, the coefficient $g \mapsto \lambda(gv_{a_o})$ is square-integrable. Consider the Haar measure on G such that Γ_{a_o} has volume 1. Then

$$\int_G |\lambda(gv_{a_o})|^2 dg = \sum_{a \in X_{n(\pi)}^1} |\lambda(v_a)|^2 < \infty ,$$

as required.

(4.1.2) **Corollary.** *If π is supercuspidal, the map $\tilde{\Psi}_\pi : (\tilde{V}, \pi^\vee) \longrightarrow \mathcal{H}_c(X_{n(\pi)}, \mathbb{C})$ induces a non-zero (whence injective) map $\bar{\Psi}_\pi : (\tilde{V}, \pi^\vee) \longrightarrow H_c^1(X_{n(\pi)}, \mathbb{C})$.*

Remark. Note that the irreducible representations of G are isomorphic to their contradredient representations.

The map $\bar{\Psi}_\pi$ is

$$\lambda \mapsto \tilde{P}si_\pi(\lambda) \bmod dC_c^0(X_{n(\pi)}, \mathbb{C}) .$$

It suffices to prove that

$$\mathcal{H}_c(X_{n(\pi)}, \mathbb{C}) \cap dC_c^0(X_{n(\pi)}, \mathbb{C}) = \{0\} .$$

If f lies in the intersection then $d^*f = 0$ and $f = dg$ for some $g \in C_c^0(X_{n(\pi)}, \mathbb{C})$. We then have $d^*\bar{f} = 0$ (where $\bar{f}(a)$ is the complex conjugate of $f(a)$), and

$$\sum_{a \in X_{n(\pi)}^1} |f(a)|^2 = \langle \bar{f}, f \rangle = \langle \bar{f}, dg \rangle = \langle d^*\bar{f}, g \rangle = 0 .$$

Hence $f = 0$.

5. The geodesic Radon transform

5.1. An *oriented apartment* $\tilde{A}$ of X is by definition a pair (A, ϵ) , where A is an apartment of X and ϵ is an orientation of A as a simplicial complex. Our group G acts on oriented apartments via $g.(A, \epsilon) = (gA, g\epsilon)$, where $g\epsilon$ is the unique orientation on A satisfying $[ga : gs]_{g\epsilon} = [a : s]_\epsilon$, for all $a \in A^1$, $s \in A^0$.

Let T be the diagonal torus of G (the image of the diagonal torus of $\mathrm{GL}(2, F)$ in G). Then the G -set $\tilde{\mathcal{A}}$ of oriented apartments in X is isomorphic to G/T . We endow $\tilde{\mathcal{A}}$ with the topology corresponding to the quotient topology of G/T . In particular for any $\tilde{A}$ in $\tilde{\mathcal{A}}$ and any open (compact) subgroup K of G , $K.\tilde{A} = \{k\tilde{A}; k \in K\}$ is an open (compact) neighbourhood of $\tilde{A}$ in $\tilde{\mathcal{A}}$. Let $\tilde{A} \in \tilde{\mathcal{A}}$ and k be a non-negative integer. By definition the (oriented) apartment of $\tilde{X}_k$ corresponding to $\tilde{A}$ is the 1-dimensional subsimplicial complex $\tilde{A}_k$ of $\tilde{X}_k$ whose edges (resp. vertices) are the $(k+1)$ -paths a (resp. k -paths s) in X contained in $\tilde{A}$ and such that the orientations of a (resp. s) and $\tilde{A}$ are compatible. We denote by $\tilde{\mathcal{A}}_k$ the set of apartments of $\tilde{A}_k$. Then $\tilde{A} \mapsto \tilde{A}_k$ is a G -equivariant bijection between $\tilde{\mathcal{A}}$ and $\tilde{\mathcal{A}}_k$ which allows us to identify both G -sets.

Let k be a non-negative integer and a be an edge of $\tilde{X}_k$. Define a subset $\tilde{\mathcal{A}}_a$ of $\tilde{\mathcal{A}}$ by

$$\tilde{\mathcal{A}}_a = \{\tilde{A} \in \tilde{\mathcal{A}}; a \in \tilde{A}^1\}.$$

(5.1.1) **Lemma** *i) With the notation as above, we have $\tilde{\mathcal{A}}_a = \Gamma_a \tilde{\mathcal{A}}_o$, for any $\tilde{\mathcal{A}}_o$ in $\tilde{\mathcal{A}}_a$. In particular $\tilde{\mathcal{A}}_a$ is a compact open subset of $\tilde{\mathcal{A}}$.*

ii) The set $\{\tilde{\mathcal{A}}_a; k \geq 0, a \in \tilde{X}_k^1\}$ is a basis of the topology of $\tilde{\mathcal{A}}$ formed of compact open subsets.

The equality $\tilde{\mathcal{A}}_a = \Gamma_a \tilde{\mathcal{A}}_o$ follows from the fact that Γ_a acts transitively on the apartments of X containing the path a . We must prove that any open subset Ω of $\tilde{\mathcal{A}}$ contains $\tilde{\mathcal{A}}_a$ for some $k \geq 0$ and $a \in \tilde{X}_k^1$. Replacing Ω by $g\Omega$ for some $g \in G$ we may assume that it contains the oriented apartment $\tilde{A}_{\mathrm{st}}$ corresponding to the coset $1.T \in G/T$. For $r \geq 1$, let K_r be the image in G of the following congruence subgroup of $\mathrm{GL}(2, F)$:

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix}; a, d \in 1 + \mathfrak{p}^r, b, c \in \mathfrak{p}^r \right\}.$$

Take r large enough so that $K_r \tilde{A}_{\mathrm{st}} \subset \Omega$. Let T^o be the maximal compact open subgroup of T . It stabilizes $\tilde{A}_{\mathrm{st}}$ pointwise, whence it

fixes $\tilde{A}_{\text{st}}$. So $K_r T^o \tilde{A}_{\text{st}} \subset \Omega$. The subgroup $K_r T^o$ is Γ_a for some $a \in \tilde{A}_{\text{st}}^1$ and we are done.

5.2. Fix $k \geq 0$. We define a (geodesic) Radon transform

$$\mathcal{R} = \mathcal{R}_k : C_c^1(\tilde{X}_k, \mathbb{C}) \longrightarrow \mathcal{F}(\tilde{\mathcal{A}}) ,$$

where $\mathcal{F}(\tilde{\mathcal{A}})$ is the space of functions on $\tilde{\mathcal{A}}$, by

$$\mathcal{R}(\omega)(\tilde{A}) = \sum_{a \in \tilde{A}^1} \omega(a) .$$

Note that the image of a 1-cochain ω whose support is reduced to a single edge a_o is $\omega(a_o) \text{Char}_{\tilde{\mathcal{A}}_{a_o}}$, where Char denotes a characteristic function. So the image of $\mathcal{R}$ actually lies in the space $C_c^0(\tilde{\mathcal{A}})$ of locally constant functions with compact support on $\tilde{\mathcal{A}}$. Clearly $\mathcal{R}$ is G -equivariant.

(5.2.1) **Lemma** *For all $f \in C_c^0(\tilde{X}_k, \mathbb{C})$, we have $\mathcal{R}(df) = 0$.*

Indeed if $\tilde{A} \in \tilde{\mathcal{A}}$, we have

$$\begin{aligned} \mathcal{R}(df)(\tilde{A}) &= \sum_{a \in \tilde{A}^1} f(a^+) - f(a^-) \\ &= \sum_{a \in \tilde{A}^1} f(a^+) - \sum_{a \in \tilde{A}^1} f(a^-) = \sum_{s \in \tilde{A}^0} f(s) - \sum_{s \in \tilde{A}^0} f(s) = 0 , \end{aligned}$$

for the map $\tilde{A}^1 \longrightarrow \tilde{A}^0$, $a \mapsto a^+$ (resp. $a \mapsto a^-$) is a bijection.

(5.2.2) **Proposition.** *a) The following sequence of G -modules is exact:*

$$C_c^0(\tilde{X}_k, \mathbb{C}) \xrightarrow{d} C_c^1(\tilde{X}_k, \mathbb{C}) \xrightarrow{\mathcal{R}} C_c^0(\tilde{\mathcal{A}}) .$$

In other words $\mathcal{R}$ induces an injective map:

$$\bar{\mathcal{R}} : H_c^1(\tilde{X}_k, \mathbb{C}) \longrightarrow C_c^0(\tilde{\mathcal{A}}) .$$

b) Moreover we have:

$$\bigcup_{k \geq 0} \bar{R}_k(H_c^1(\tilde{X}_k, \mathbb{C})) = C_c^0(\tilde{\mathcal{A}}) .$$

Point b) follows from the fact that $C_c^0(\tilde{\mathcal{A}})$ is generated as a $\mathbb{C}$ -vector space by the functions $\text{Char}_{\tilde{\mathcal{A}}_a}$, where $a \in \tilde{X}_k^1$ and $k \geq 0$.

To prove a) we need to introduce some more concepts. A path p in $\tilde{X}_k$ is a sequence of edges a_u , $u = 0, \dots, l-1$, such that for $u = 0, \dots, l-2$, a_u and a_{u+1} share a vertex. Abusing the notation we shall write $p = (x_0, x_1, \dots, x_l)$, where for $u = 0, \dots, l-1$, $\{a_u^+, a_u^-\} = \{x_u, x_{u+1}\}$, keeping in mind that when $k = 0$ two neighbour vertices do not determine a unique edge. We define incidence coefficients $[p : a_u] \in \{\pm 1\}$, $u = 0, \dots, l-1$, by $[p : a_u] = 1$ if and only if $a_u^- = x_u$ and $a_u^+ = x_{u+1}$.

Let $\omega \in C_c^1(\tilde{X}_k, \mathbb{C})$ be in the kernel of $\mathcal{R}$. The “integral” of ω along p is by definition

$$\int_p \omega = \sum_{u=0, \dots, l-1} [p : a_u] \omega(a_u) .$$

(5.2.3) **Lemma** *With the notation as above, if p is a loop, i.e. if a_0 and a_{l-1} share the vertex $x_l = x_0$, then*

$$\int_p \omega = 0 .$$

We first show that the lemma implies proposition (5.2.2). Fix $s_o \in \tilde{X}_k^0$ and $\alpha_o \in \mathbb{C}$. For any $x \in \tilde{X}_k^0$, we set

$$f(x) = \alpha_o + \int_p \omega ,$$

where $p = (x_0, \dots, x_l)$ is any path satisfying $x_0 = s_o$ and $x_l = x$. We claim that $f(x)$ does not depend on the choice of p . Indeed let $q = (y_0, \dots, y_m)$ be another path satisfying the same assumptions and set $q - p = (z_0, \dots, z_{m+l+1})$, where $z_u = x_u$, for $u = 0, \dots, l$, and $z_u = y_{m+l+1-u}$, for $u = l+1, \dots, l+1+m$. Then one easily checks that

$$(\alpha_o + \int_q \omega) - (\alpha_o + \int_p \omega) = \int_{q-p} \omega = 0 ,$$

since $q - p$ is a loop.

Let $a \in \tilde{X}_k^1$ and $p = (x_o, \dots, x_l)$ be a path such that $x_0 = s_o$ and $x_l = a^-$. Set $q = (x_o, \dots, x_l, a^+)$. Then

$$f(a+) - f(a-) = \int_q \omega - \int_p \omega = [q : a] \omega(a) = \omega(a) .$$

So we must now prove that one can choose s_o and α_o so that $f \in C^0(\tilde{X}_k, \mathbb{C})$ has finite support. Let S_k be the support of ω in $\tilde{X}_k^1$ and set

$$S := \bigcup_{a \in S_k} \text{cvx}(a) \subset X ,$$

where $\text{cvx}(a)$ denotes the convex hull of a in (the geometric realization of) X . Then S is a bounded subset of X . Let t be a vertex in S and δ be an integer large enough so that $S \subset X(t, \delta)$, where $X(t, \delta)$ is the subtree of X whose vertices are at combinatorial distance from t less than or equal to δ . Then the complementary set ${}^cX(t, \delta)$ of $X(t, \delta)$ in X has the following property; for any k -path $a \subset {}^cX(t, \delta)$, there exists a half-apartment A^+ such that $a \subset A^+ \subset {}^cX(t, \delta)$. Set

$$S'_k = \{s \in \tilde{X}_k^0 ; \text{cvx}(s) \cap X(t, \delta) \neq \emptyset\}.$$

Choose s_o outside S'_k and set $\alpha_o = 0$. We are going to prove that the support of f is contained in S'_k . Let x be in $\tilde{X}_k^0 \setminus S'_k$. Choose half-apartments A_o^+ and A^+ so that:

$$A^+, A_o^+ \subset {}^cS \text{ and } \text{cvx}(s_o) \subset A_o^+, \text{cvx}(x) \subset A^+ .$$

Consider vertices s'_o and x' of $\tilde{X}_k$ whose convex hulls in X lie in A_o^+ and A^+ respectively. One may choose s'_o (resp. x') away enough from the origin of A_o^+ (resp. of A^+) so that there exists an apartment B of X containing both $\text{cvx}(s'_o)$ and $\text{cvx}(x')$.

First case: the vertices s'_o and x' induce the same orientation on B (see figure 1). Let $\tilde{B}$ be the corresponding oriented apartment. Let $\tilde{A}_o^+$ be the oriented half-apartment whose orientation is induced by s_o and $\tilde{A}^+$ be the oriented half-apartment whose orientation is induced by x . Let $p(s_o, s'_o)$ (resp. $p(x', x)$ and $p(s'_o, x')$) be the unique injective path in $\tilde{X}_k$ joining s_o to s'_o (resp. x' to x , s'_o to x') and such that $p(s_o, s'_o) \subset \tilde{A}_o^+$ (resp. $p(x', x) \subset \tilde{A}^+$, $p(s'_o, x') \subset \tilde{B}$). By concatenation, we get a path $p(s_o, x) = p(s_o, s'_o) + p(s'_o, x') + p(x', x)$ joining s_o and x . Since those vertices of $\tilde{X}_k$ occuring in $p(s_o, s'_o)$ or $p(x', x)$ do not lie in S , we have

$$\int_{p(s_o, s'_o)} \omega = \int_{p(x, x')} \omega = 0 .$$

Moreover

$$\int_{p(s'_o, x')} \omega = \pm \mathcal{R}(\omega)(\tilde{B}) = 0 ,$$

by assumption. So

$$\int_{p(s_o, x)} \omega = 0 \text{ and } f(x) = \alpha_o + \int_{p(s_o, x)} \omega = 0 ,$$

as required.




Figure 1

Second case: the vertices s'_o and x' induce different orientations on B (see figure 2). Then one can choose a third vertex $x'' \in S'_k$ such that s'_o and x'' (resp. x' and x'') lie in some commun apartment B_1 (resp. B_2) and induce the same orientation on that apartment. We denote by $\tilde{B}_1$ and $\tilde{B}_2$ the corresponding oriented apartments. Then, with the notation as in the first case, one easily shows that

$$f(x) = f(s_o) + \int_{p(s_o, s'_o)} \omega + \pm \mathcal{R}(\omega)(\tilde{B}_1) + \pm \mathcal{R}(\omega)(\tilde{B}_2) + \int_{p(x', x)} \omega = 0 .$$




Figure 2

Proof of lemma (5.2.3). It is based on the following easy lemma whose proof is left to the reader.

(5.2.4) **Lemma.** *Let U be either $\mathbb{Z}$, $\mathbb{N}$ or a finite interval of integers. Let $p = (x_u)_{u \in U}$ be a path in $\tilde{X}_k$. Assume that p satisfies one of the following properties:*

- (P1) *For all $u \in U$ such that $u + 1 \in U$, we have $a_u^+ = a_{u+1}^-$;*
- (P2) *For all $u \in U$ such that $u + 1 \in U$, we have $a_u^- = a_{u+1}^+$.*

Then p is injective and there is an apartment $\tilde{A}$ containing p . In particular p cannot be a loop.

Remark. A path p satisfies (P1) or (P2) if and only if the sequence of incidence numbers $([p : a_u])_u$ is constant.

Let $p = (x_0, \dots, x_l)$ be a loop. We consider the index u as an element of $\mathbb{Z}/l\mathbb{Z}$. According to the previous lemma, the set V of indices $u \in \mathbb{Z}/l\mathbb{Z}$ such that we have neither $a_u^+ = a_{u+1}^-$ nor $a_u^- = a_{u+1}^+$ is non-empty. Moreover if it is non-empty it has cardinal at least 2. Let us first consider the case $\sharp V = 2$ (this case indeed occurs when $k = 0$). We may for instance assume that

$$a_0^- = a_{l-1}^- = x_0 \text{ and } a_{u_o-1}^+ = a_{u_o}^+ = x_{u_o} ,$$

for some $u_o \in \mathbb{Z}/l\mathbb{Z} \setminus \{0\}$. Hence we must have

$$a_{u+1}^- = a_u^+, \quad u = 0, \dots, u_o - 2 \quad a_u^- = a_{u+1}^+, \quad u = u_o, \dots, l - 2 .$$




Figure 3

Choose a half-apartment A_o^+ , whose vertex set is $\{s_0, s_1, \dots, s_u, \dots\}$, satisfying:

- for all $u \geq 0$, there is an edge b_u of A_o^+ , such that $b_u^+ = s_u$, $b_u^- = s_{u+1}$;
- $s_0 = x_0 = b_0^+$.

Similarly choose a half-apartment $A_{u_o}^+$, whose vertex set is $\{t_0, t_1, \dots, t_u, \dots\}$, satisfying:

- for all $u \geq 0$, there is an edge c_u of $A_{u_o}^+$ such that $c_u^+ = s_{u+1}$, $c_u^- = s_u$;
- $t_0 = x_{u_o} = c_0^-$.

Consider the two infinite paths:

$$p_1 = (\dots, s_u, \dots, s_1, s_0 = x_0, x_1, x_2, \dots, x_{u_o-1}, x_{u_o} = t_0, t_1, t_2, \dots, t_v, \dots)$$

$$p_2 = (\dots, s_u, \dots, s_1, s_0 = x_0, x_{l-1}, x_{l-2}, \dots, x_{u_o+1}, x_{u_o} = t_0, t_1, t_2, \dots, t_v, \dots)$$

By lemma (5.2.4) we can find an apartment $\tilde{A}_1$ (resp. $\tilde{A}_2$) whose vertices are those of p_1 (resp. those of p_2). Since ω has finite support, we can give an obvious meaning to the integrals:

$$\int_{p_1} \omega \text{ and } \int_{p_2} \omega .$$

Moreover we have

$$\int_{p_1} \omega = \mathcal{R}(\omega)(\tilde{A}_1) = 0 \text{ and } \int_{p_2} \omega = \mathcal{R}(\omega)(\tilde{A}_2) = 0 .$$

Finally we have:

$$\begin{aligned}
\int_{p_1} \omega - \int_{p_2} \omega &= \sum_{u=0, \dots, u_o-1} \omega(a_u) - \sum_{u=l-1, l-2, \dots, u_o} \omega(a_u) \\
&= \sum_{u=0, \dots, u_o-1} [p : a_u] \omega(a_u) + \sum_{u=l-1, l-2, \dots, u_o} [p : a_u] \omega(a_u) = \int_p \omega = 0 ,
\end{aligned}$$

QED.

This proof extends to the case $\#V > 2$ by introducing for all $u \in V$ a half-apartment starting at the vertex $a_u \cap a_{u+1}$. The details are left to the reader (see figure 4).




Figure 4

6. Some uniqueness results

6.1. Uniqueness of the model in $H_c^1(\tilde{X}_k, \mathbb{C})$ for supercuspidals. We may rephrase proposition (5.2.2) in the following way.

(6.1.1) **Proposition.** *For all $k \geq 0$, the Radon transform induces an injective G -equivariant map:*

$$H_c^1(\tilde{X}_k, \mathbb{C}) \longrightarrow c\text{-ind}_T^G 1_T ,$$

where $c\text{-ind}$ denotes a compactly induced representation and 1_T denotes the trivial character of T .

(6.1.2) **Theorem** *Let $(\pi, \mathcal{V}_\pi)$ be an irreducible supercuspidal representation of G . Then we have*

$$\dim_{\mathbb{C}} \text{Hom}_G[\mathcal{V}_\pi, H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})] = 1 .$$

This theorem indeed follows from the following result due to Waldspurger ([8] Prop. 9', p. 31):

(6.1.3) **Theorem** (J.-L. Waldspurger). *Let $(\pi, \mathcal{V}_\pi)$ be an irreducible unitary smooth representation of G . Then we have*

$$\dim_{\mathbb{C}} \text{Hom}_G(\mathcal{V}_\pi, 1_T) = 1 .$$

However we shall not use this result. We are going to show how the uniqueness of Whittaker model (see e.g. [5] Theorem 2.14) implies that

$$\dim_{\mathbb{C}} \text{Hom}_G(\mathcal{V}_\pi, H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})) \leq 1$$

for all irreducible supercuspidal representations of G .

Let U be the unipotent radical of the upper triangular Borel subgroup of G . We use the notation

$$u(t) := \begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix} \in U, \quad t \in F .$$

We fix a non-trivial smooth character ψ_F of $(F, +)$ and set $\theta(u(t)) = \psi_F(t)$, $t \in F$. Attached to the non-degenerate character θ of U we have the space of θ -Whittaker functions $\text{Wh}_\theta = \text{Ind}_U^G \theta$. Here Ind denotes a smooth induced representation. We shall use the convention that functions in Wh_θ transforms according to θ under the right action of U and that G acts on the left.

(6.1.4) **Proposition.** *i) For all $f \in c\text{-ind}_T^G 1_T$ and $g \in G$, the function $f_g : F \rightarrow \mathbb{C}$, $t \mapsto f(gu(t))$ is locally constant and has compact support.*

ii) For all $f \in c\text{-ind}_T^G 1_T$, set

$$W_f(g) = \int_U f(gu(t)) \bar{\psi}_F(t) dt .$$

Then $\Phi : c\text{-ind}_T^G 1_T \rightarrow \text{Wh}_\theta$, $f \mapsto W_f$ is an injective G -equivariant map.

So theorem (6.1.2) follows now from proposition (6.1.1) and the uniqueness of the Whittaker model of an irreducible smooth representation $(\pi, \mathcal{V}_\pi)$ of G :

$$\dim_{\mathbb{C}}(\mathcal{V}_\pi, \text{Ind}_U^G \theta) \leq 1 .$$

Proof of (6.2.4). i) Let $f \in c\text{-ind}_T^G 1_T$ and $g \in G$. Since f is smooth, f_g is smooth as well. There exists a compact subset C of G such that the support of f is contained in CT . So for a $\lambda \in F$ such that $f(gu(\lambda)) \neq 0$, there exists $t_\lambda \in T$ such that $u(\lambda)t_\lambda \in g^{-1}C$. Since the product map $F \times T \longrightarrow UT$, $(\lambda, t) \mapsto u(\lambda)t$ is a homeomorphism, the set of $\lambda \in F$ such that $f(gu(\lambda)) \neq 0$ is contained in a compact subset of F .

ii) With the notation of the proposition, the map Φ is clearly G -equivariant. Let $f \in \ker \Phi$. The functions f_g , $g \in G$ belong to the Schwartz-Bruhat space of locally constant functions on F with compact support. We are going to prove that their Fourier transforms $\hat{f}_g$, $g \in G$, are zero. This will implies that $f = 0$. Recall that

$$\hat{f}_g(\mu) = \int_F f(gu(\lambda)) \bar{\psi}_F(\mu\lambda) d\lambda, \mu \in F, g \in G .$$

By a change of variable we get:

$$\hat{f}_g(\mu) = \frac{1}{|\mu|_F} \int_F f(gu(\mu^{-1}\lambda)) \bar{\psi}_F(\lambda) d\lambda, \mu \in F, \mu \neq 0, g \in G .$$

For $\mu \neq 0$, choose $t_\mu \in T$ such that $t_\mu u(\lambda) t_\mu^{-1} = u(\mu^{-1}\lambda)$. Since f is right T -invariant, for all $g \in G$, we get:

$$\hat{f}_g(\mu) = \frac{1}{|\mu|_F} \int_F f(gt_\mu u(\lambda)) \bar{\psi}_F(\lambda) d\lambda = \frac{1}{|\mu|_F} \Phi(f)(gt_\mu) = 0 .$$

So for all $g \in G$, $\hat{f}_g$ vanishes on $F \setminus \{0\}$, whence must be zero since it is locally constant.

6.2. *A new proof of a result of Casselman.* Let $(\pi, \mathcal{V}_\pi)$ be an irreducible supercuspidal representation of G . Recall that in §3 we defined $n(\pi) \in \mathbb{N}$ by

$$\mathcal{V}^{\Gamma \circ (\mathfrak{p}^{n(\pi)+1})} \neq \{0\} \text{ and } \mathcal{V}^{\Gamma \circ (\mathfrak{p}^{n(\pi)})} = \{0\} .$$

We give a new proof of the uniqueness part of Casselman's theorem (3.1.3) based on proposition (6.1.1)

(6.2.1) **Proposition.** *With the notation as above, we have*

$$\dim_{\mathbb{C}} \mathcal{V}^{\Gamma_o(\mathfrak{p}^{n(\pi)+1})} = 1 .$$

Let v_o, w_o be two non-zero vectors in $\mathcal{V}^{\Gamma_o(\mathfrak{p}^{n(\pi)+1})}$. Using the construction of §3, we attach to v_o and w_o two injective and G -equivariant maps $\tilde{\Psi}^{v_o}, \tilde{\Psi}^{w_o} : \tilde{\mathcal{V}} \longrightarrow H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$. They are the compound maps of the injection $\mathcal{H}_c(\tilde{X}_{n(\pi)}, \mathbb{C}) \longrightarrow H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$ and of $\tilde{\Psi}^{v_o}, \tilde{\Psi}^{w_o} : \tilde{\mathcal{V}} \longrightarrow \mathcal{H}_c(\tilde{X}_{n(\pi)}, \mathbb{C})$. The latter maps are given by $\tilde{\Psi}^{v_o}(f)(a) = f(gv_o)$, $\tilde{\Psi}^{w_o}(f)(a) = f(gw_o)$, for all $f \in \tilde{\mathcal{V}}, a \in \tilde{X}_{n(\pi)}^1, a = ga_o, g \in G$. By theorem (6.2.2), there exists $\lambda \in \mathbb{C}^\times$ such that $\tilde{\Psi}^{v_o}(f) = \lambda \tilde{\Psi}^{w_o}(f)$, $f \in \tilde{\mathcal{V}}$. So for all $g \in G, f \in \mathcal{V}^\vee$, we have $f(gv_o) = \lambda f(gw_o)$, i.e. $f(g(v_o - \lambda w_o)) = 0$. If $v_o - \lambda w_o \neq 0$, then the $g(v_o - \lambda w_o), g \in G$, generate $\mathcal{V}$ and any $f \in \tilde{\mathcal{V}}$ must be zero: a contradiction. So $v_o = \lambda w_o$ as required.

References

- [1] P. Broussous, *Simplicial complexes lying equivariantly over the affine building of $GL(N)$* , Mathematische Annalen, **329**, 2004, 495–511.
- [2] G. Ahumada Bustamante, *Analyse harmonique sur l'espace des chemins d'un arbre*, PhD thesis, University of Paris Sud, 1988.
- [3] P. Cartier, *Harmonic analysis on trees*, Harmonic Analysis on Homogeneous Spaces, Calvin C. Moore, ed., Proc. Sympos. Pure Math., **XXVI**, Amer. Math. Soc., Providence, R. I., 1973, 419–424.
- [4] W. Casselman, *On some result of Atkin and Lehner*, Math. Ann. **201**, 1873, 301–314.
- [5] H. Jacquet, R.P. Langlands, *Automorphic forms on $GL(2)$* , Lecture Notes in Math., **114**, Springer Verlag, 1970.
- [6] H. Jacquet, I. Piatetski-Shapiro, J. Shalika, *Conducteur des représentations du groupe linéaire*, Math. Ann., **256** no. 2, 1981, 199–214.
- [7] J.-P. Serre, *Trees*, Springer, 2nd ed. 2002.
- [8] J.-L. Waldspurger, *Correspondance de Shimura*, J. Math. Pures et Appl., **59**, 1980, 1–133.